

2-2021

Evaluating the Efficacy of Telestroke Intervention in a Large Community Hospital Network

Daniel Moylan

Daniel Joffe

Charles Morse

Michael Knapp

Ahmad Sweid

See next page for additional authors

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2023_phase1



Part of the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Daniel Moylan, Daniel Joffe, Charles Morse, Michael Knapp, Ahmad Sweid, and Pascal Jabbour

Evaluating the Efficacy of Telestroke Intervention in a Large Community Hospital Network

Daniel Moylan, Daniel Joffe**, Charles Morse**, Michael Knapp, Ahmad Sweid*, Pascal Jabbour

Disclosures: None

Introduction

Literature Review:

Telemedicine for Acute Ischemic Stroke (Telestroke) allows live consultations with remote specialists¹

IV-tPA for AIS has significant benefits, but only when administered within 3-4.5 hours of onset²⁻⁴

In previous studies, telestroke has been shown to be safe and to improve rates of thrombolysis in smaller patient populations^{4,5}

TIME IS BRAIN



Importance:

Acute Ischemic Stroke Accounts for 87% of acute strokes and represents a significant *global* burden diminishing QOL^{6,7}

Only 3.4-5.2% of AIS patients in the US receive thrombolytic therapy due to difficulties in rapid recognition and delivery⁸

The efficacy of telestroke intervention with regards to patient outcomes is not well described in larger studies

Mechanical Thrombectomy⁹





Objectives & Hypothesis

Research Question

How does telestroke intervention compare to previous standard of care with regards to IV-tPA administration and patient outcomes for AIS patients in a large, multicenter network of community hospitals?

Hypothesis

Access to telestroke care across the Thomas Jefferson University Hospital Network will result in increased thrombolytic perfusion rates and improved outcomes for acute ischemic stroke patients.

Approach

Design: Retrospective cohort study

Population: 9,702 patients evaluated for acute ischemic stroke at 36 Thomas Jefferson Hospitals from 2014-2019

Intervention: Implementation of Telestroke robot system across Thomas Jefferson Hospital Network

Data Source: TJUH Telestroke database, medical records, literature (control)

Rationale: Utilizes existing data – RCT not ethical since IV-tPA administration benefits are well documented

Data Collection (Completed)

Demographics – Sex, ethnicity, age, hospital, department, vessel

Thrombolysis – Administration (Y/N), indications, contraindications, time of onset to consult, time of onset to administration

Patient Outcomes – NIHSS Stroke scale, mortality, MT performance & complications, discharge destination

Results - Outcomes

Preliminary Data Analysis

- Large sample size + COVID Issues have delayed complete data analysis


IV-tPA Administration Rate

- 3.8-5.2% in literature vs. 8.3% in the experiment group
- Expected to increase

NIHSS Stroke Scale Outcome

- IV-tPA administration associated with a significant improvement in NIHSS stroke scale ($p < 0.0001$; 95% confidence interval [CI] = 4.27, 7.80)

IV - tPA Administration Rates



Conclusions

- Telestroke is a new, affordable, rapidly expanding technology that has not yet been tested on a broad scale
 - Coronavirus has highlighted the importance of remote care
- Telestroke intervention for AIS is feasible, effective, and safe in a large hospital network
 - Rates of IV-tPA administration are increased when compared to current rates in literature
 - Significant improvements in NIHSS Stroke scale associated with IV-tPA
- Impact: Data supports the implementation of telestroke systems on a similarly large scale
 - Improved accessibility to stroke care, particularly in underserved areas

Future Directions



Prospective study design

- Study limited by retrospective and literature controls
- May be possible at new site testing or implementing telestroke

Expansion of mobile stroke unit + telestroke intervention

- Ambulance that is equipped for stroke care
- Quicker access to stroke care
- Allows areas without any stroke center to receive care

Acknowledgements

All fellow authors + IT Support (Carlos Desarden)

References

- 1) Demaerschalk BM, Berg J, Chong BW, et al. American Telemedicine Association: Telestroke Guidelines. *Telemed J E Health*. 2017;23(5):376–389.
- 2) Lattimore SU, Chalela J, Davis L, et al. Impact of establishing a primary stroke center at a community hospital on the use of thrombolytic therapy: the NINDS Suburban Hospital Stroke Center experience. *Stroke*. 2003;34(6):e55-7
- 3) Campbell BC. Thrombolysis and Thrombectomy for Acute Ischemic Stroke: Strengths and Synergies. *Semin Thromb Hemost*. 2017;43(2):185-190.
- 4) Madhavan M, Karceski S. Telestroke: Is it safe and effective? *Neurology*. 2016;87(13):e146-8.
- 5) Legris N, Hervieu-Begue M, Daubail B, et al. Telemedicine for the acute management of stroke in Burgundy, France: an evaluation of effectiveness and safety. *Eur J Neurol*. 2016;23(9):1433-40.
- 6) Mozaffarian D, Benjamin EJ, Go AS, et al. Heart Disease and stroke statistics—2015 update: a report from the American Heart Association. *Circulation*. 2015;131(4):e29-322.
- 7) Murray CJ, Lopez AD. Measuring the global burden of disease. *N Engl J Med*. 2013;369(5):448-57.
- 8) Adeoye O, Hornung R, Khatri P, Kleindorfer D. Recombinant tissue-type plasminogen activator use for ischemic stroke in the United States: a doubling of treatment rates over the course of 5 years. *Stroke*. 2011;42(7):1952–1955.
- 9) Meng SW, Kuo RC, Yang HJ, Lai CL, Wu CC, Hsieh MY. Recruiting an Acute Coronary Team to Perform Emergent Mechanical Thrombectomy in Acute Ischemic Stroke Patients: A Successful Case and Team Model in a Local Hospital. *Acta Cardiol Sin*. 2018 Jan;34(1):99-103.



Jefferson

Philadelphia University +
Thomas Jefferson University

HOME OF SIDNEY KIMMEL MEDICAL COLLEGE